

Sarion

- Sarion has a lower nitrogen requirement
- High Phytophthora resistance
- High starch content



Characteristics

| | |
|------------------------------|----------------------------------|
| Variety | Sarion |
| Maturity | Late |
| Tuber – Shape | Round |
| Tuber – Flesh colour | Light yellow |
| Tuber – Skin colour | Yellow |
| Yield | High |
| Destination | Starch processing |
| Blight – Foliage | 8,5 |
| Blight – Tuber | 7 |
| Common scab | 6,5 |
| Leaf roll | 7 |
| A - Virus | 7 |
| X - Virus | 8 |
| Yn - Virus | 8 |
| Drought | 8,5 |
| Nematode – Ro 1,4 | 9 / rs 0,0 % |
| Nematode – Ro 2,3 | 9 / rs 0,0 % |
| Nematode – Pa 2 | 9 / rs 0,0 % |
| Nematode – Pa 3 | 9 / rs 0,0 % |
| Nematode tolerance | 6,5 |
| Wart disease - Fysio 1 | Resistant |
| Wart disease - Race 2/6 | 8 |
| Wart disease - Race 18 | Resistant |
| Fertilization Advice N | 140-180 kg/ha (incl. soil stock) |
| Fertilization Advice P | Based on soil analysis |
| Fertilization Advice K | 250-280 kg/ha |
| Planting distance (35/55 mm) | 30 - 33 cm |